

- Mota J, Dogan S, Brkić J, Teixeira JP, Valdiglesias V. Medication use in older patients and age-blind approach: narrative literature review (insufficient evidence on the efficacy and safety of drugs in older age, frequent use of PIMs and polypharmacy, and underuse of highly beneficial nonpharmacological strategies). *European journal of clinical pharmacology*. 2019; 4: 1–6.
3. Sørbye LW, Garms-Homolová V, Henrard JC, Jónsson PV, Fialová D, Topinková E, Gambassi G. Shaping home care in Europe: the contribution of the Aged in Home Care project. *Maturitas*. 2009; 62(3): 235–242.
4. Onder G, Liperoti R, Fialová D, Topinková E, Tosato M, Danese P, Gallo PF, Carpenter I, Finne-Soveri H, Gindin J, Bernabei R, Landi F; Shelter Project. Polypharmacy in nursing home in Europe: results from the Shelter study. *J Gerontol A Biol Sci Med Sci*. 2012; 67(6): 698–704.
5. Fialová D, Topinková E, Gambassi G, Finne-Soveri H, Jónsson PV, Carpenter I, Schroll M, Onder G, Sørbye LW, Wagner C, Reissigová J, Bernabei R, ADHOC project research group. Potentially inappropriate medication use among elderly home care patients in Europe. *JAMA*. 2005; 293 (11): 1348–1358.
6. Fialová D. The RAI-MDS (SHELTER, ADHOC) medication datasets and their use for evaluating appropriateness of medication prescribing. In: *Aging Clinical and Experimental Research* (2nd ed.) 2011; 23(1): 47 (ISSN 1720-8319). VIth International Association of Gerontology and Geriatrics (IAGG) – European Region Congress „Healthy and Active Ageing for All Europeans II“, Bologna, Italy.
7. Lix LM, Yan L, Blackburn D, HU N, Schneider-Lindner V, Shevchuk Y, Teare GF. Agreement between administrative data and the Resident Assessment Instrument Minimum Dataset (RAI-MDS) for medication use in long-term care facilities: a population-based study. *BMC geriatrics*. 2015; 15: 24. doi:10.1186/s12877-015-0023-2.
8. Morris JN, Fries BE, Morris SA. Scalling ADLs within the MDS. *The Journals of Gerontology. Series A: Biological Sciences and Medical Sciences*. 1999; 54(11): M546–M553. doi:10.1093/gerona/54. 11. m546
9. Morris JN, Fries BE, Mehr DR, et al. The MDS Cognitive Performance Scale. *J Gerontol*. 1994; 49 (4): 174–182.
10. Burrows AB, Morris JN, Simon SE, Hirdes JP, Phillips CH. Development of a Minimum Data Set-based depression rating scale for use in nursing homes. *Age Aging*. 2000; 29: 165–172.
11. Somers AI, Mallet L, Van der Cammen T, Robays H, Petrovic M. Applicability of an adapted medication appropriateness index for detection of drug-related problems in geriatric inpatients. *Am J Geriatr Pharmacother*. 2012; 10(2): 101–109.
12. Curtin D, Gallagher PF, O'mahony D. Explicit criteria as clinical tools to minimize inappropriate medication use and its consequences. *Therapeutic advances in drug safety*. 2019; 10: 1–10. ISSN 2042-0986.
13. Fick DM, Cooper JW, WADE WE, WALLER JL, MACLEAN JR, BEERS MH. Updating the Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. *Arch Intern Med*. 2003; 163: 2716–2724.
14. American geriatrics society 2012 beers criteria update expert panel. The American Geriatrics Society 2012 Beers Criteria Update Expert Panel. *J Am Geriatr Soc*. 2012; 60(4): 616–631.
15. Barry PJ, O'keefe N, O'connor KA, O'mahony D. Inappropriate prescribing in the elderly: a comparison of the Beers criteria and the improved prescribing in the elderly tool (IPET) in acutely ill elderly hospitalized patients. *J Clin Pharm Ther*. 2006; 31(6): 617–626.
16. Gallagher P, Ryan C, Byrne S, Kennedy J, O'mahony D. Stoppp (Screening Tool of Older Persons' Prescriptions) and Start (Screening Tool to Alert Doctors to Right Treatment): Consensus Validation. *Int J Clin Pharm Ther*. 2008; 46(2): 72–83.
17. Fialová D, Topinková E, Ballóková A, Matějovská Kubešová H. Expertní konsensus ČR 2012 v oblasti léčiv a lékových postupů potenciálně nevhodných ve stáří. Oddíl I. Vhodnost volby léčiv a dávkovacích schémat u geriatrických pacientů Oddíl II Interakce lék–nemoc ve stáří. (2012 CZ expert consensus for potentially inappropriate medication use in old age: Appropriate choice of drugs and drug dosing in geriatric patients (Section I), drug-disease interactions in the old age (Section II) *Klinická farmakologie a farmacie*. 2013; 27(1): 18–28.
18. R core team. R: A Language and Environment for Statistical Computing. [software] version 3. 4. 2. Vienna, Austria: R Foundation for Statistical Computing, 2017-09-28. Dostupné z: <https://www.R-project.org>.
19. Davidoff AJ, Miller GE, Sarpong EM, Yang E, Brandt N, Fick DM. Prevalence of Potentially Inappropriate Medication Use in Older Adults Using the 2012 Beers Criteria. *Journal of the American Geriatrics Society*. 2015; 63(3): 486–500.
20. Hudhra K, García-Caballeros M, Jucja B, Casado-Fernández E, Espigares-Rodríguez E, Bueno-Cavanillas A. Frequency of potentially inappropriate prescriptions in older people at discharge according to Beers and STOPP criteria. *International Journal of Clinical Pharmacy*. 2014; 36(3): 596–603. ISSN 2210-7703.
21. Fialová D, Brkić J, Laffon B, Reissigová J, Grešáková S, Dogan S, Doro P, Tasić L, Marinković V, Valdiglesias V, Costa S, Kostřba J. Applicability of EU (7)-PIM criteria in cross-national studies in European countries. *Ther Adv Drug Saf*. 2019; 10: 2042098619854014.
22. Gallagher P, Lang PO, Cherubini A, Topinková E, Cruz-Jentoft A, Montero Errasquín B, Mádllová P, Gasperini B, Baeyens H, Baeyens JP, Michel JP, O'mahony D. Prevalence of potentially inappropriate prescribing in an acutely ill population of older patients admitted to six European hospitals. *European Journal of Clinical Pharmacology*. 2011; 67(11): 1175–1188.
23. San-José A, Agustí A, Vidal X, Formiga F, Gómez-hernández M, García J, López-Soto A, Ramírez-Duque N, Torres OH, Barbé J. Potentially Inappropriate Prescription in Older Patients in Spain (PIPOPS) Investigators' project. Inappropriate prescribing to the oldest old patients admitted to hospital: prevalence, most frequently used medicines, and associated factors. *BMC Geriatr*. 2015; 15: 42.
24. Ballóková A, Peel NM, Fialová D, Scott IA, Gray LC, Hubbard RE. Use of Benzodiazepines and Association with Falls in Older People Admitted to Hospital: A Prospective Cohort Study. *Drugs and aging*. 2014; 31(4): 299–310.
25. Zelinková A. Analýzy racionality užívání hypnotik u seniorů v akutní péči v českém souboru projektu EUROAGEISM H2020 (dipl.práce, školitel: doc. D. Fialová, obhájena na FaF UK červen 2019). (online) Dostupná z: https://is.cuni.cz/studium/dipl_uc/index.php?id=65d4f29ab714a15cdd3757454b9a1ba6&tid=1&do=main&doo=detail&did=194368 (poslední přístup: 29. 8. 2020).
26. Puldová K. Stárnutí populace a specifické aspekty lékové preskripce hypnotik ve stáří (II.) (dipl.práce, školitel: doc. D. Fialová, obhájena na FaF UK červen 2020). (online) Dostupná z: https://is.cuni.cz/studium/dipl_uc/index.php?id=65d4f29ab714a15cdd3757454b9a1ba6&tid=2&do=main&doo=detail&did=205741 (poslední přístup: 29. 8. 2020).
27. Pek EA, Remfry A, Pendrith C, Fan-Lun C, BHATIA RS a SOONG C. High Prevalence of Inappropriate Benzodiazepine and Sedative Hypnotic Prescriptions among Hospitalized Older Adults. *Journal of Hospital Medicine*. 2017; 12(5): 310–316.
28. Diem SJ, Ewing SK, Stone KL, Ancoli-Israel S, Redline S, Ensrud KE. Use of Non-Benzodiazepine Sedative Hypnotics and Risk of Falls in Older Men. *Journal of Gerontology*. 2014; 3(3): 158. ISSN 21677182.
29. Ballóková-Lukačšínová A, Fialová D. Benzodiazepines, age-related pharmacological changes, and risks in older adults. In: Preedy VR. *Neuropathology of Drug Addictions and Substance Misuse. Volume 3: General Processes and Mechanisms, Prescription Medications, Caffeine, Polydrug Misuse, Emerging Addictions and Non-Drug Addictions*. 2016, Elsevier Inc., pp: 334–344 (ISBN 978-0-12-800634-4).
30. Masaryková P. Riziká gastrotoxicity a racionalita užitia inhibitorov protónovej pumpy u seniorov v akútnej starostlivosti v projekte EUROAGEISM H2020 (dipl.práce, školitel: doc. D. Fialová, obhájena na FaF UK červen 2019). (online) Dostupná z: https://is.cuni.cz/studium/dipl_uc/index.php?id=65d4f29ab714a15cdd3757454b9a1ba6&tid=3&do=xdownload&fid=120348496&did=194367&vdetailu=1
31. Marroquín CE, Inglesia MN a Cobos PL. Adequacy of medication in patients 65 years or older in teaching health centers in Cáceres, Spain. *Revista Española de Salud Pública*. 2012; 86(4): 419–434.
32. Mukai Y, TAMPI RR. Treatment of depression in the elderly: A review of the recent literature on the efficacy of single-versus dual-action antidepressants. *Clinical Therapeutics*. 2009; 31(5): 945–961.
33. Wawruch M, Fialová D, Zikavská M, et al. Factors influencing the use of potentially inappropriate medication in older patients in Slovakia. *Journal of Clinical Pharmacy and Therapeutics*. 2008; 33(4): 381–392.
34. Leendertse AJ, Van Den Bemt PM, Poolman JB, Stoker LJ, Egberts AC, Postma MJ. Preventable hospital admissions related to medication (HARM): cost analysis of the HARM study. *Value Health*. 2011; 14(1): 34–40.