

**Tab. 1.** Vliv fenobarbitalu na farmakokinetiku léků z jiných indikačních skupin (10)

| | hladina | AUC | CL | t _{1/2} |
|---|-----------|--------|---------|------------------|
| 9-aminokamptothecin | ↓ 67 % | | ↑ 68 % | |
| albendazol* | ↓ 63 % | ↓ 71 % | | ↓ 39 % |
| atenolol | | ↓ 24 % | | |
| cyklosporin | ↓ 95 % | | | |
| desipramin | ↓ 30 % | | | |
| dextropropoxyfen | ↑ 20 % | | | |
| diazepam N-desmethyldiazepam | ↑ | | ↑ 158 % | ↓ 62 % |
| etoposid | | | ↑ 77 % | ↓ 18 % |
| felodipin | ↓ 82 % | ↓ 94 % | | |
| glufosfamid | | ↓ 15 % | ↑ 15 % | |
| chinidin | | | ↑ 60 % | ↓ 50 % |
| chloramfenikol | ↓ 70–95 % | | | |
| irinotekan [†] 7-ethyl-10-hydroxykamptothecin | ↓ 75 % | | | |
| lidokain | | ↓ 60 % | ↑ 196 % | |
| metotrexát | | | ↑ 55 % | |
| metronidazol | | ↓ 33 % | ↑ 50 % | |
| nifedipin | | ↓ 60 % | | |
| nimodipin | | ↓ 86 % | | |
| paclitaxel | | ↓ 52 % | ↑ 101 % | |
| paracetamol [‡] | | ↓ 39 % | ↑ 52 % | |