

HLAVNÍ TÉMA

SOUČASNÉ MOŽNOSTI OPERAČNÍHO ŘEŠENÍ KRČNÍ SPONDYLOGENNÍ MYELOPATIE

5. Breig A, Turnbull I, Hassler O. Effects of mechanical stresses on the spinal cord in cervical spondylosis. A study on fresh cadaver material. *J Neurosurg.* 1966;25:45-56.
6. Cloward RB. The anterior approach for removal of ruptured cervical discs. *J Neurosurg.* 1958;15:602-614.
7. Fehlings MG, Skaf G. A review of the pathophysiology of cervical spondylotic myelopathy with insights for potential novel mechanisms drawn from traumatic spinal cord injury. *Spine.* 1998;23:2730-2737.
8. Fehlings M, Barry S, Kopjar B, Yoon ST, et al. Anterior versus posterior surgical approaches to treat cervical spondylotic myelopathy: outcomes of the prospective multicenter AOSpine North America CSM study in 264 patients. *Spine (Phila Pa 1976).* 2013;38(26):2247-52.
9. Fehlings MG, Wilson JR, Karadimas SK, et al. Clinical evaluation of a neuroprotective drug in patients with cervical spondylotic myelopathy undergoing surgical treatment: design and rationale for the CSM-protect trial. *Spine.* 2013;38:68-75.
10. Fehlings MG, Tetreault LA, Riew KD, et al. A Clinical Practice Guideline for the Management of Patients With Degenerative Cervical Myelopathy: Recommendations for Patients With Mild, Moderate, and Severe Disease and Nonmyelopathic Patients With Evidence of Cord Compression. *Global Spine J.* 2017;7(3 Suppl):70-83.
11. Hashimoto M, Mochizuki M, Aiba A, et al. C 5 palsy following anterior decompression and spinal fusion for cervical degenerative diseases. *Eur Spine J.* 2010;19:1702-1710.
12. Hirabayashi K, Watanabe K, Wakano K, et al. Expansive open-door laminoplasty for cervical stenotic myelopathy. *Spine.* 1983;8:693-699.
13. Jiang L, Chen W, Chen Q, et al. Clinical application of a new plate fixation system in open-door laminoplasty. *Orthopedics.* 2012;35(2):225-231.
14. Kadaňka Z, Benešová Y, Mechl M. Krční myelopatie – diagnostický souhrn. *Česk Slov Neurol N.* 2009;72/105(2):110-114.
15. Kadaňka Z. Spondylogenní cervikální myelopatie. *Česk Slov Neurol N.* 2010;73/106(3):209-226.
16. Kato S, Nouri A, Wu D, et al. Comparison of Anterior and Posterior Surgery for Degenerative Cervical Myelopathy. *JBJS Surgery.* 2017;99(12):1013-1021.
17. Kovalova I, Kerkovsky M, Kadanka Z, et al. Prevalence and imaging characteristics of asymptomatic and symptomatic spondylotic cervical spinal cord compression. *Spine.* 2016;41(24):1908-1916.
18. Kurokawa T, Tsuyama N, Tanaka H, et al. Enlargement of the spinal canal by the sagittal splitting of the spinous process. *Bessatsu Seikeigeka.* 1982;2:234-240.
19. Lin S, Zhou F, Sun Y, et al. The severity of operative invasion to the posterior muscular-ligament complex influences cervical sagittal balance after open-door laminoplasty. *Eur Spine J.* 2015;24:127-135.
20. Luo J, Cao K, Huang S, et al. Comparison of anterior approach versus posterior approach for the treatment of multilevel cervical spondylotic myelopathy. *Eur Spine J.* 2015;24:1621-1630.
21. Matsunaga S, Komiya S, Toyama Y. Risk factors for development of myelopathy in patients with cervical spondylotic cord compression. *Eur Spine J.* 2015;24:142-149.
22. McCormick JR, Sama AJ, Schiller NC, Butler AJ, Donnelly CJ 3rd. Cervical Spondylotic Myelopathy: A Guide to Diagnosis and Management. *J Am Board Fam Med.* 2020;33(2):303-313.
23. Mesfin A, Park MS, Piyaskulkaew Ch, Chuntarapas T, et al. Neck pain following laminoplasty. *Global Spine J.* 2015;5:17-22.
24. Nurick S. The natural history and the results of surgical treatment of the spinal cord disorder associated with cervical spondylosis. *Brain: J Neurol.* 1972;95(1):101-108.
25. Paracino R, Fasinella MR, Mancini F, et al. Review of laminoplasty versus laminectomy in the surgical management of cervical spondylotic myelopathy. *Surg Neurol Int.* 2021;12:44.
26. Ryba L, Chaloupka R, Repko M, Cienciala J. Dlouhodobé výsledky operačního řešení cervikální spondylogenní myelopatie pomocí „open-door“ laminoplastiky. *Acta Chir Orthop Traumatol Cech.* 2015;3:209-215.
27. Sakara H, Hosono N, Mukai Y, et al. C5 palsy after decompression surgery for cervical myelopathy: review of the literature. *Spine (Phila Pa 1976).* 2003;28:2447-2451.
28. Sakaura H, Hosono N, Mukai Y, et al. C3-6 laminoplasty for cervical spondylotic myelopathy maintains satisfactory long-term surgical outcomes. *Global Spine J.* 2014;4:169-174.
29. Schmeiser G, Bergmann JJ, Papavero L, Kothe R. Surgical Treatment of Multilevel Degenerative Cervical Myelopathy: Open-Door Laminoplasty and Fixation via Unilateral Approach. A Feasibility Study. *J Neurol Surg A Cent Eur Neurosurg.* 2021. doi: 10.1055/s-0041-1739224.
30. Smith GW, Robinson RA. The treatment of certain cervical spine disorders by the anterior removal of the intervertebral disc and interbody fusion. *J Bone Joint Surg.* 1958;40 A:607-624.

5. NEURO-MUSKULÁRNÍ FÓRUM

NEUROMUSKULÁRNÍ SEKCE ČNS



21.–22. 9. 2023

Hotel Kraskov
Třemošnice-Starý Dvůr



VÍCE INFORMACÍ NA
www.kongresneurologie.cz

